				2614				
RABIES CERTIFICAT	ΓΕ							
Species: Cat Weight: Under 20 lbs Sex:	□ Male, altere	d □ Fe	male, alt	ered				
Age: □ 3-5 mo old □ 6–12 mo old □ ≥ 12 mo old	old Color: _							
Breed:	nted							
Producer: Fort Dodge RabVac 3 Serial #:								
Other vaccines: Fort Dodge Fel-O-Vax LVK III (FeLV, FPV,								
Veterinarian signature:	, , - ,							
Veterinarian name & license #:								
OPERATION CATNIP MEDICA								
	Amount	Route	Time	Initials				
Anesthesia TKX	mL	IM	Tillic	IIIIIIIII				
TKX 2 nd dose (if needed)	mL	IM						
Microchip scan No chip Chip #								
Antibiotic Dual penicillin (right front leg) 1.0 m	nL 0.5 mL	SC						
Ear tipping Left								
Vaccines FVRCP/FeLV (LHL) Rabies (RHL)		SC						
Analgesia Buprenorphine (oral transmucosal) 0.1 m	nL 0.05 mL	OTM						
Parasiticide Adv. Multi (Imidacloprid & Moxidectin) 0.23mL	L 0.4mL 0.8mL	Topical						
Fluids Lactated Ringers 0.9% Saline	mL	SC						
Reversal Yohimbine (medial saphenous)	mL	IV						
Other								
OPERATION CATNIP SURGICA	AI DECODO							
	Veterinarian Veterinary student Full name: Injectable only Isoflurane% by mask Other:							
Spay approach Midline Left flank Other:								
• Ovarian ligatures 2-0 3-0 Suture type:								
Uterine body ligatures 2-0 3-0 Suture type:								
Body wall closure 2-0 3-0 Suture type:	2-0 3-0 Suture type: Continuous Interrupted							
• Subcutaneous closure 2-0 3-0 Suture type:	Continu		terrupted					
 Skin closure Condition 2-0 3-0 Suture type: Routine Pregnant # fetus 	None ses In heat		dhesive actating					
• Condition Routine Pregnant # fetus Neuter approach Scrotal Abdominal Inguinal	ses iii iieat	Lo	acialing					
Cord ligation Open Closed Autoligation								
• Condition Routine Cryptorchid: L-Inguinal		Abdominal	R-Abdon	 ninal				
Already neutered Requires Dr. Levy/DVM Supervisor co	•	ls prior to cl	osing incis	ion:				
Other findings URI Abscess Other:								
Other treatments								
Notes								
SPECIAL EXAM REQUI	IRFD							
Surgeon to check								
Outcome of exam								
NOTES TO CAREGIVER:								

				• Date: @gmail.com •						
				RABIES CERT						
Species:	Cat	Weig	ı ht: Undeı	20 lbs S e	ex: 🗆 M	lale, altere	ed □ Fe	male, alt	ered	
Age:	□ 3-5	mo old	□ 6–12	mo old □ ≥ 1	2 mo old	Color:				
Breed:		H □ DI	mo old □ 6–12 mo old □ ≥ 12 mo old Color: 							
Producer				erial #:						
				ax LVK III (FeL\						
	ian signa		010.00	ax 2 v · · · · · (· · o 2	·, · ,	.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	- milai c		2000:01	
	•									
vetermar	ian name	& licens		TON CATNID M						
		Tv		TION CATNIP M procedures)	Ame		Route Time Initials			
An	esthesia	TKX	pe (circle)	nocedules)	Alli	mL	IM	Tillic	IIIIIIIII	
7 111	ootiiiooia		lose (if need	ded)		mL	IM			
Microc	hip scan	No chip	Chip #	•						
Α	ntibiotic	Dual peni	cillin (right f	ront leg)	1.0 mL	0.5 mL	SC			
Ea	r tipping	Left								
Vaccines FVRCP/F			Rabies (RHL)			SC				
		phine (oral transmucosal)		0.1 mL		OTM				
		ti (Imidacloprid & Moxidectin)		0.23mL 0.4		Topical				
			Ringers 0.9% Saline			mL	SC			
				e (medial saphenous)		mL	IV			
	Other									
			OPERATI	ON CATNIP SU	JRGICAL I	RECORD				
Anasthas	sia		Injectable	only leoflurane	% by m	ack Othe	or.			
Anesthesia Spay approach		Injectable only Isoflurane% by mask Other: Midline Left flank Other:								
Ovarian ligatures		2-0 3-0 Suture type: Autoligation								
	ne body liga		2-0 3-0 Suture type:							
					Continuous Interrupted					
Subcutaneous closureSkin closure		2-0 3-0 2-0 3-0	Suture type: Suture type:		Continu None		terrupted dhesive			
• Skin	Ciosuie		2-0 3-0	Suture type.		None		unesive		
Neuter a	proach		Scrotal	Abdominal In	guinal					
	Cord ligation Open Closed Autoligation									
				Inguinal R-Inguinal L-Abdominal R-Abdominal						
Already neuteredRequires Dr. Levy/DVM SupOther findingsURIAbscessOth				mation/initia	als prior to cl	osing incis	ion:			
Other tre	_		URI	Abscess Oth	er:					
Notes	atments									
.10103			<u> </u>							
			SI	PECIAL EXAM	REQUIRE	D				
Surgeon to check										
Outcome of exam										
NOTES TO	CAREGIVER:									

Surgery Recovery Instructions



Thank you for bringing stray or feral cats for sterilization today. Your concern has helped lessen the burden of cat overpopulation. The cats were given a long acting injection of anesthetic and pain medication for today's surgery. They may still be under the influence of these drugs at the time of discharge, even if they appear to be fully recovered. Consequently, you should monitor the cats for a minimum of 24 hours

following surgery in order to provide the best care possible for the cats.

PLEASE READ THE FOLLOWING CAREFULLY:

- Please keep all cats in their traps until the day after surgery, even if they appear to be
 friendly and alert. This provides protection and security for both the cats and for the people
 that work with them.
- The cats will be groggy and need to sleep overnight. Please do not disturb them by reaching into the trap they often overreact and may bite! Anyone who is bitten or scratched by a cat should receive immediate medical attention. Do not release a cat that has caused an injury until authorized by Alachua County Animal Services.
- Do not give adult cats food or water in the trap the same day as surgery was performed.
 They may spill the water and may vomit and aspirate if fed. You can give them food and water the next morning when they are fully awake.
- **Kittens should be offered a small amount of canned food** the night of surgery as soon as they are fully recovered from anesthesia. This helps keep their blood sugar normal.
- Keep the cats in their traps in a warm, draft-free area like a bathroom or garage until the
 next morning. Cats should never be left outside following surgery. The cats cannot regulate
 their body temperature while recovering from anesthesia and are susceptible to death from
 heat or cold exposure.
- Normal behaviors that occur during recovery include head bobbing, wobbly movements, agitation, shivering, and hypersensitivity to light and sound. No cat should be released the next day if it continues to exhibit these signs.
- **Abnormal behaviors during recovery** include continued bleeding from the surgery incisions, difficulty breathing, vomiting, and inability to awaken.
- If the cats are fully awake the next day, they may be released back to their colony and offered food and water. If the cat is not yet fully awake or is bleeding, please call the Operation Catnip emergency number.

If you have any concerns about the recovery of the cats, please call the emergency hotline at (352) 258-6658 to discuss the situation with a volunteer on-call veterinarian. Non-emergency questions should be directed to the main Operation Catnip phone number (352) 380-0940 or email at ocgainesville@gmail.com during regular business hours. If a cat has a situation that you feel cannot wait for assistance from Operation Catnip, please call your local pet emergency clinic. Payment of fees for care at a veterinary clinic is your responsibility, not Operation Catnip's.

CAT LOVERS LIKE YOU make Operation Catnip possible. Please support our efforts by joining the Catnip volunteer team. Even with the generosity of so many volunteers, it costs Operation Catnip about \$30 to help each cat. **\$30 gives a life-saving and life-changing chance** to a kitty who has spent every day scrounging and trying to make it on her own. Please give as generously as you can.